## APPENDIX

ANTERIOR KNEE PAIN (Sheet code:	)
Name:	_ Date:
Age:	
Knee: L/R	
Duration of symptoms: years months	
For each question, circle the latest choice (letter), which corresp	oonds to your knee symptoms.
<ul> <li>1. Limp <ul> <li>(a) None (5)</li> <li>(b) Slight or periodical (3)</li> <li>(c) Constant (0)</li> </ul> </li> <li>2. Support <ul> <li>(c) Support</li> </ul> </li> </ul>	<ul> <li>8. Prolonged sitting with the</li> <li>(a) No difficulty (10)</li> <li>(b) Pain after exercise (8)</li> <li>(c) Constant pain (6)</li> <li>(d) Pain forces to extend knees</li> <li>(e) Unable (0)</li> </ul>
<ul> <li>(a) Full support without pain (5)</li> <li>(b) Painful (3)</li> <li>(c) Weight bearing impossible (0)</li> <li><b>3. Walking</b></li> <li>(a) Unlimited (5)</li> <li>(b) More than 2 km (3)</li> <li>(c) 1-2 km (2)</li> <li>(d) Unable (0)</li> </ul>	<ul> <li>9. Pain <ul> <li>(a) None (10)</li> <li>(b) Slight and occasional (8)</li> <li>(c) Interferes with sleep (6)</li> <li>(d) Occasionally severe (3)</li> <li>(e) Constant and severe (0)</li> </ul> </li> <li>10. Swelling <ul> <li>(a) None (10)</li> </ul> </li> </ul>
<ul> <li>4. Stairs <ul> <li>(a) No difficulty (10)</li> <li>(b) Slight pain when descending (8)</li> <li>(c) Pain both when descending and ascending (5)</li> <li>(d) Unable (0)</li> </ul> </li> </ul>	<ul> <li>(b) After severe exertion (8)</li> <li>(c) After daily activities (6)</li> <li>(d) Every evening (4)</li> <li>(e) Constant (0)</li> </ul>
<ul> <li>5. Squatting</li> <li>(a) No difficulty (5)</li> <li>(b) Repeated squatting painful (4)</li> <li>(c) Painful each time (3)</li> <li>(d) Possible with partial weight bearing (2)</li> <li>(e) Unable (0)</li> </ul>	<ul> <li>11. Abnormal painful kneeca (subluxations)</li> <li>(a) None (10)</li> <li>(b) Occasionally in sports activitie</li> <li>(c) Occasionally in daily activitie</li> <li>(d) At least one documented di</li> <li>(e) More than two dislocations</li> </ul>
<ul> <li>6. Running</li> <li>(a) No difficulty (10)</li> <li>(b) Pain after more than 2 km (8)</li> <li>(c) Slight pain from start (6)</li> <li>(d) Severe pain (3)</li> </ul>	<b>12. Atrophy of thigh</b> (a) None (5) (b) Slight (3) (c) Severe (0)
<ul> <li>(e) Unable (0)</li> <li><b>7. Jumping</b> <ul> <li>(a) No difficulty (10)</li> <li>(b) Slight difficulty (7)</li> <li>(c) Constant pain (2)</li> <li>(d) Unable (0)</li> </ul> </li> </ul>	<ul> <li>13. Flexion deficiency</li> <li>(a) None (5)</li> <li>(b) Slight (3)</li> <li>(c) Severe (0)</li> </ul>
Reference: Kujala UM, Jaakkola LH, Koskinen SK, Taim patellofemoral disorders. <i>Arthroscopy</i> 1993, <b>9:</b> 159-163	

knees flexed

temporarily (4)

## ap (patellar) movements

- ities (6) es (4) lislocation (2) (0)