



MANAGING PAIN SAFELY

Pain is an expected part of having surgery.

Managing your pain safely is important to help your recovery.

PAIN AFTER SURGERY



After surgery, a moderate amount of pain is a normal and expected part of the process.

The goal of pain medication is to keep you moving and help you cope with your pain, but these medications *will not eliminate your pain entirely*.

In general, the most painful time is the first 72 hours (3 days) after surgery.

The amount of pain usually decreases after these first few days.



Scan to watch a short video on managing your pain

MANAGING PAIN SAFELY

Opioids, such as morphine, oxycodone (Percocet), tramadol, hydromorphone or codeine, are strong pain relief medications that may be prescribed to help you manage the most intense postoperative pain.

Opioids should be used with caution as research shows that up to 1 in 12 patients prescribed an opioid for short-term use after surgery are at risk of becoming chronic users.

The regular use of over-the-counter Tylenol® and Advil, in addition to Cold Therapy*, can effectively manage pain and significantly reduce, or eliminate the need for opioids.

If you are still experiencing intolerable pain after regularly using Tylenol® and Advil in addition to Cold Therapy*, take the opioid in the lowest dose possible.

Also, only use the opioid for the shortest amount of time. No more than 2 - 7 days.



*Scan for more information on Cold Therapy



Banff Sport Medicine

CONTACT US

If you have questions about managing your pain, or your pain increases or does not decrease with medication, please contact our team:

- Banff Sport Medicine 403-760-2897 ext 1 (during business hours 8 am - 4 pm Monday to Friday)
- Banff Mineral Springs Hospital 403-762-2222 (for urgent concerns outside of regular business hours)





Banff Sport Medicine

Opioid Medication: Pain Control after Surgery

What are my options for safe and effective pain control?

Managing your pain after surgery is important. Combination therapy (using different medications together) along with education can offer the best pain relief. Some of these medications work via different pathways and can be used together. After surgery a moderate amount of pain is a normal and expected part of the process. Pain medications help you function better and cope with the amount of pain you are experiencing, but **these medications will not eliminate your pain entirely.**

Talk to your surgeon about medication options, which can include both prescription strength and over the counter drugs. These medications include:

- **Non-steroidal Anti-inflammatory Drugs (NSAIDs)** can be used to decrease swelling and fever, and to treat mild to moderate pain. (These can be prescription or over-the-counter medications, such as Naprosyn, naproxen: *Aleve*, ibuprofen: *Advil*, *Motrin*).
- **Acetaminophen** is used to decrease mild to moderate pain and fever. (These are over-the-counter medications such as Tylenol or Paracetamol).
- **Local anesthetics** which can be injected near a set of nerves (nerve block) or the surgery wound site. (This option is performed immediately after your surgery at the hospital).
- **Opioids** are used to decrease severe pain. (These are prescription medications such as Morphine, Tramacet, Percocet, OxyContin, Vicodin or Tylenol #3 which contains codeine).
- **Nerve pain medication** used to prevent nerve pain and irritation. (These are prescription medications such as Lyrica and Neurontin).

When should I take pain medication?

The goal is to keep you moving and control your pain. In general, the most painful time is the first 72 hours (3 days) after surgery. The amount of pain usually decreases after these first few days. If your pain increases or does not decrease with medication you should contact your surgeon. Call our office at 403-760-2897 (during regular business hours 8am-4pm Monday to Friday), or the Banff Mineral Springs Hospital at 403-762-2222 (outside of business hours).

Some medication is taken on a schedule (for example, every 4-6 hours) and other medications can be taken when you feel pain. Here is a useful guide:

- I am moving slowly and either have no pain or only a little pain = I don't need any medication.
- I am feeling some pain and having trouble moving around = I could use some over-the-counter medication. (For example, ibuprofen: *Advil/Motrin* or acetaminophen: *Tylenol*).
- I am thinking about my pain all the time and it is painful for me to move = I should take some stronger medication. (For example, a prescription medication such as Morphine, Percocet or Tramacet).

It usually takes 25-40 minutes for the medications to start working effectively so you should take your medication before your pain becomes severe or you are unable to move because of the pain.

Are there ways to reduce my pain besides medication?

Yes! You can do a number of things to help decrease your pain (*continued on next page*)

- Use ice packs or cold therapy to decrease swelling and pain.
- Rest for the majority of time for the first few days after surgery.
- Elevate and support your surgical limb as shown on your rehabilitation exercise protocol.
- Keep moving after the first few days, without over doing it.

- Distraction such as watching movies, listening to music, playing games, or talking to friends and family, has been shown to help reduce pain after surgery.

Why are opioids used?

Opioids are strong medications and when used properly, they can help to relieve short-term pain like the pain experienced after surgery. When opioids are used improperly, they can cause dependence or addiction, overdose and death.

How to use your prescribed opioid medication.

If you have been prescribed an opioid medication, such as Percocet or Tramacet, it should:

- **Only** be taken as prescribed
- **Never** be used by someone for whom it was not prescribed
- **Never** be taken with alcohol or other medications (except as prescribed).
- **Never** to be used while driving or operating machinery

Keep your medication safe to help prevent use by others by:

- **Never** sharing your medication with anyone else. (Sharing opioid medication is illegal and may also cause serious harm or death to the other person).
- **Keeping track** of the number of pills remaining in the package
- **Storing** opioids in a safe and secure place, out of the reach of children and teenagers

Unused portions of opioid medicine should always be:

- Kept out of sight of children and pets
- Stored in a safe place to prevent theft, problematic use or accidental exposure
- **Returned to the pharmacy** for safe disposal when no longer needed or expired. Medications should never be thrown in the trash or flushed down the toilet.
- This will prevent the possibility of illegal use and protect the environment from contamination.

Opioid Side Effects

Short-term side effects	Long-term side effects
Drowsiness	Increased tolerance
Decreased reaction time, which can impair driving and decision-making	Substance use disorder or dependence (addiction)
Constipation	Liver damage
Impotence in men	Infertility in women
Nausea and vomiting	Worsening pain
Difficulty breathing, which can lead to or worsen sleep apnea	Life-threatening withdrawal symptoms in babies born to mothers taking opioids
Euphoria (feeling high)	Overdose
Headaches, dizziness and confusion, which can lead to falls or fractures	

Anyone who takes prescription opioids can become addicted.

If you take opioid medications for more than a few weeks, your body becomes used to that dose. This physical dependence means you may experience withdrawal symptoms if you stop taking the drug.

Withdrawal symptoms include: insomnia, anxiety, racing heartbeat, and headaches. Withdrawal symptoms can be managed by gradually decreasing your medication dose with advice from a health care provider.

Signs of opioid overdose include small pupils, trouble breathing and unconsciousness. Call 911 immediately if you think you or a family member took too much pain medication.